


AD A 045965

**INJURIES AND ILLNESSES OF VIETNAM WAR POWS.
II. ARMY POWS**

S. W. BERG & M. RICHLIN

REPORT NO. 75-82



AD No. _____
DDC FILE COPY



✓ **NAVAL HEALTH RESEARCH CENTER**

SAN DIEGO, CALIFORNIA 92152

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND

BETHESDA, MARYLAND

Injuries and Illnesses of Vietnam War POWs. II. Army POWs

LCDR S. William Berg, MC, USN*
Milton Richlin, Ph.D.

THE first report in this series¹ presented the results of repatriation medical examinations of US Navy personnel who had been held as prisoners of war (POWs) in Vietnam. The present report describes the medical status of the 77 repatriated POWs (RPWs) who were members of the US Army.

The characteristics of the Army RPWs may be summarized as follows: there were 28 officers (including 10 Warrant Officers) and 49 enlisted men; the average age at time of capture was 27.6 years for officers, and 23.0 years for enlisted men; duration of captivity (in months) averaged 44.3 for officers, and 52.1 for enlisted men.

Duty assignments at time of capture included: 23 helicopter pilots and aircrew members, 16 combat infantry, 10 Special Forces, eight advisors, seven truck drivers, two fixed-wing aircrewmembers, and 11 miscellaneous.

The present report describes the procedures and findings of the medical teams that examined and treated the RPWs at Clark Air Force Base. (Republic of the Philippines), and at eight mainland Army hospitals.

All Army RPWs had been captured in South Vietnam (SVN), and spent several weeks to several years there, involuntarily sharing the harsh life of a roving guerilla band.²⁻³ The majority of the RPWs were transferred, after varying periods of time, to enter the prison system in North Vietnam (NVN), but 23 per cent spent their entire captivity in the South.

A brief overview of prison conditions can serve as a backdrop for understanding the physical and psychological stresses this group of men endured during captivity. In SVN, POWs were held in small groups accompanying guerilla bands as they travelled in the mountain jungle or wet, swampy areas.²⁻³ Shelter consisted of bamboo and thatch "hootches" or sometimes bunkers dug in the soil. If the area was sufficiently far from American troops and the jungle or swamp sufficiently difficult to travel through, the men would not be restrained; otherwise, they were chained or manacled inside their cages. In addition to having to endure whatever extremes of weather that were present, particularly the monsoon rains which flooded the hootches, the POWs lived in an endemic malaria area. They shared the guerillas' fate of being at the tail end of the supply chain, which usually meant inadequate amounts of food, medicine, and other vital supplies. Rice was eaten almost exclusively, although sometimes manioc (a tuberous plant similar

to a yam) was substituted. The POWs also ate whatever greens, fish, small animals, or reptiles they could scavenge.³⁻⁵

Medical care for POWs held in SVN was much more limited than in NVN. Although in some secure areas the Viet Cong had set up field hospitals, no American POWs were treated there. One Army physician was held captive in SVN for a few years, but he was not allowed to treat his fellow POWs (although he did so covertly).³ Medical care was provided by the guerillas themselves, who sometimes had a soldier attached to their unit who functioned as a corpsman. The quantity of medicines available (including vitamin B₁ for beriberi and quinine for malaria) was always extremely limited. POWs rarely received medication until they were desperately ill, due, in part, to the fact that use of medicines to treat a POW might mean that none would be available if one of the guerillas fell ill.

In NVN, POWs were held in conventional prisons, or buildings that had been converted to prisons, in or near metropolitan areas.* Not only was food better quantitatively and qualitatively in NVN than in SVN but, after late 1969, the quantity of food increased, and conditions improved generally in NVN, but not in SVN.

In the present report, the emphasis is on documentation of the diagnoses made at "Operation Homecoming." Where appropriate, information is also presented concerning symptoms and conditions which occurred during captivity (as described in histories obtained by Operation Homecoming physicians).

Methods

The repatriation plan, originally drawn up under the code name "Egress Recap," was later given the name "Operation Homecoming." Details of the repatriation plan at "Operation Homecoming" (O/H) were described previously.¹ Briefly, all RPWs were admitted to the sicklist for a minimum of 90 days during O/H, with most RPWs spending the first two to four weeks actually in the hospital. During this period, the RPW was in the care of a primary physician, usually an internist, although general medical officers and surgeons were also utilized. All RPWs were evaluated by other consultants as needed.

O/H directives called for recording of all positive and negative findings. The basic tool to accomplish this documentation was the Initial Medical Evaluation Form (IMEF), a 400-page, 29-section instrument detailing all procedures to be done and providing space to record the results. The details of this document have been described in our earlier report.¹

After each RPW was discharged from the sick list, his complete O/H medical record was forwarded to the Center

From the Center for Prisoner of War Studies, Naval Health Research Center, San Diego, Calif. 92152.

*Present address, Dept. of Medicine, Naval Regional Medical Center, San Diego, Calif. 92134.

This research was supported by the Department of the Navy, Bureau of Medicine and Surgery, under Work Order Request Number N00018-75-WR-00004 dated 1 July 1974 and by the Office of the Surgeon General, Department of the Army under Military Interdepartmental Purchase Request Number 75-01 dated 23 September 1974.

*Details of the captivity experience in North Vietnam have been described previously.^{1,4,6}

Table 1
Most Common Diagnoses Among Army Vietnam RPWs at Operation Homecoming

ICDA*	Diagnosis	Freq.	Percent
120.0-129.9	Helminthiases	59	77
260.0-269.9	Avitaminoses & Other Nutritional Deficiencies	42	55
389.9	Hearing Impairment	31	40
111.0-110.9	Dermatophytosis	30	39
952.0-957.9	Peripheral Nerve Injury	30	39
084.0-084.9	Malaria	26	34
370.0-370.9	Refractive Errors	22	29
307.0	Transient Situational Disturbance	19	25
377.0-377.9	Other Diseases of Retina & Optic Nerve	18	23
706.0-706.9	Disease of Sebaceous Gland	17	22
40000-40009**	Amoebiasis	17	22
460.0-466.9	Acute Upper Respiratory Infection	14	18
805.0-805.9	Fractured Vertebrae (Compression)	14	18
891.0-891.9	Open Wound - Knee, Leg, Ankle	13	17
782.6-787.2	Edema - Swelling	12	16
454.0	Hemorrhoids	11	14
680.0-686.9	Skin Infections (Bacterial)	11	14
733.6	Residual Foreign Body	11	14
728.0-728.9	Vertebrogenic Pain Syndrome	10	13
377.2	Amblyopia, Acquired	9	12
054.0	Herpes Simplex	8	10
079.1	Viral Warts	8	10
692.9	Eczema & Dermatitis	8	10
757.1	Pigmented Nevus	8	10
713.1	Spondylitis Osteoarthritis	7	9
873.0-873.9	Laceration - Head	7	9
306.8-791.0	Cephalalgia - Headaches	6	8
360.0	Conjunctivitis	6	8
504.0	Deflected Nasal Septum	6	8
507.0	Hay Fever	6	8
519.2	Inactive Pulmonary Granulomatous Disease	6	8
872.9	Open Wound - Ear, Late Effect	6	8
40200-40212**	Positive PPD	6	8
111.0	Pityriasis Versicolor	5	6
285.9	Anemia, Unspecified	5	6
373.0-373.9	Strabismus	5	6
470.0-474.9	Influenza	5	6
480.0-486.9	Pneumonia	5	6
714.0	Traumatic Arthritis	5	6
725.0-725.9	Schmorl's Nodules	5	6
735.0	Scoliosis	5	6
756.1	Anomalies of Lumbosacral Joint	5	6
781.3	Disturbed Hearing (Tinnitus)	5	6
945.0-945.9	Burn - Lower Limb	5	6
004.0-004.9	Bacillary Dysentery	4	5
273.5	Gilbert's Syndrome	4	5
280.0	Iron Deficiency Anemia	4	5
300.0-300.9	Neuroses	4	5
377.3	Color Blindness	4	5
490.0	Bronchitis	4	5
550.0	Inguinal Hernia	4	5
564.0-564.9	Functional Disorder - Intestine	4	5
690.0	Seborrheic Dermatitis	4	5
875.0-875.9	Open Wound - Chest	4	5
881.0-881.9	Open Wound - Elbow, Forearm, Wrist	4	5
890.0-890.9	Open Wound - Hip, Thigh	4	5

* ICDA: Code numbers taken from U.S. National Center for Health International Classification of Diseases for Use in the U.S., 8th Rev.

** Code numbers created by CPWS Medical Specialties Branch

for Prisoner of War Studies (CPWS) in San Diego for study and microfilming. Included were medical records from Clark AFB, complete hospital in-patient charts, and the IMEF forms. All diagnoses, abnormal signs, symptoms, and laboratory results were extracted and recorded on individual coding sheets, along with pertinent ancillary information relating to etiology, history, treatment (if any), and status of each diagnosis at time of the RPW's release from the sick list. Diagnostic code numbers were assigned using the *International Classification of Diseases, Adapted for United States, Eighth Edition (ICDA-8)*.† Occasionally the ICDA-8 system was inadequate (e.g., positive PPD reaction) and, in these instances, a CPWS coding system was used. All data were key-punched for computer analysis, and analyzed using standard statistical programs.

Results

The 77 RPWs accumulated 1,149 diagnoses (an average of 14.9 per RPW), which comprised 386 separate diagnostic entities.

Table 1 lists, in rank order, the most frequently occurring diagnoses. The frequency count gives the number of RPWs presenting with each particular diagnosis.‡

Among the numerous and diverse diagnoses not appearing in Table 1 due to their low frequency, several warrant mention because of their clinical significance. One RPW was severely wounded in the lumbar spine at the time of capture, cutting part of the cauda equina at the L-5 level, producing paraplegia and accompanying urinary and fecal incontinence. Another RPW was wounded in the left parietal area, with sequelae including both generalized and focal motor seizures, moderate expressive aphasia, spastic right hemiparesis, and a right-central facial neurologic defect. A third RPW, age 20, also suffered both a head wound producing persisting generalized convulsive seizures, and a left-side cerebrovascular accident which produced a five-hour loss of consciousness and a left hemiparesis of four-months duration during captivity. At the time of release, he had fully recovered from his hemiparesis. Still another RPW suffered brief amnesia and a transient left hemiparesis at the time of shutdown. A fifth RPW was severely beaten during captivity and, at repatriation, x-rays of his skull showed a calcified intracerebral hematoma.

There was one RPW with mild essential hypertension at repatriation, and two other RPWs had minor ECG abnormalities. One man was successfully resuscitated during captivity by the Army POW-flight surgeon, after suffering what was diagnosed at O/H as a pulmonary embolus.

A variety of gastrointestinal diagnoses were made at O/H, notably two duodenal ulcers, two peptic ulcers, and one case of esophagitis.

†The painstaking work of research assistants Nancie Cicalo, John Deaton, Carol Million, and Nancy Shreve is gratefully acknowledged.

‡The frequency count indicates the number of RPWs presenting with the given diagnosis. Where a number of diagnoses have been combined (e.g., Helminthiases - 120.0-129.9), an RPW is counted only once in the "category" even though he may have both Ascariasis - 127.0 and Trichuriasis - 127.2. In several cases, an RPW presented with two or more problems, each carrying the same diagnostic label; in this report the RPW is counted only once for a given diagnosis.

Five RPWs passed renal stones in captivity (as indicated by rather classic histories). These were attributed by them to dehydration due to the limited amount of drinking water provided. At repatriation, there were two diagnoses of non-venereal urethritis, two of prostatitis, and three miscellaneous urinary tract infections.

Respiratory system diagnoses at O/H included: 14 acute upper respiratory infections and four acute bronchitis. An additional four RPWs presented with influenza. On the basis of prolonged productive cough and fever, three RPWs believed that they had developed pneumonia during captivity. Chronic bronchitis was diagnosed in two RPWs by O/H physicians, largely on the basis of a history of chronic cough. Episodes resembling acute asthmatic attacks occurred in two RPWs during captivity, but have not recurred since repatriation. These attacks were of sufficient severity to prompt the captors to treat the men with unknown pills and injections.

Table 2 lists the various diagnoses of malnutrition. Additional diagnoses and historical symptoms whose etiologies may include nutritional aspects were as follows: nine diagnoses of nutritional amblyopia, mostly of minimal degree; two RPWs with radiological evidence of osteoporosis; and 31 RPWs who reported that edema or swelling of the extremities (usually of the lower legs and lasting from a few days to several weeks) had occurred during captivity.

Transient, mild laboratory abnormalities were seen in many of the RPWs. Occasionally, the BUN was elevated, generally in the range of 25 to 35, for about one to four weeks. The serum creatine was normal in these individuals, as were additional renal function studies when done. Similarly, the SGOT and SGPT, and sometimes the LDH, were mildly elevated for two to three weeks. There was one case of acute refeeding pancreatitis, although complaints of vague, low grade abdominal discomfort were common the first few days after release. This may have been due to the RPWs gorging themselves on traditional American foods long denied them and the problems were treated symptomatically. There was one case of refeeding gynecomastia. There were three cases of asymptomatic elevated serum uric acid levels, and four RPWs were diagnosed as having Gilbert's syndrome.

Table 2

Malnutrition in Army Vietnam RPWs

ICDA *	DIAGNOSIS	FREQ.	PERCENT
260	Vitamin A deficiency	5	6
261	Thiamine deficiency	12	16
263	Other B vitamin deficiency	7	9
264	Ascorbic acid deficiency	3	4
265	Vitamin D deficiency	1	1
266	Vitamin deficiency, not further specified	8	10
267	Protein malnutrition	1	1
269	Sprue	1	1
269.9	Malnutrition, not further specified	22	29
280.0-291.9	Deficiency anemia	6	8

* ICDA: Code numbers taken from U.S. National Center for Health International Classification of Diseases for Use in the U.S., 8th Rev.

Table 3 presents the infectious and parasitic diseases diagnosed at O/H. Nearly all of the RPWs were convinced that they had "worms." This was based on frequently seeing worms in their stools in captivity, or occasionally plucking an ascaris out of one's nose or mouth. Their convictions are supported by the fact that ova or cysts were found in fecal smears in 77 per cent of the RPWs at O/H (Table 1).

All the RPWs had diarrhea in captivity, especially while held in the South. The frequency and duration of diarrheal episodes did, however, vary considerably from one RPW to another. Most of the RPWs were convinced that they had dysentery in captivity, and several described watery diarrheal stools containing blood and mucus and occurring several times a day. Others used a "formula": 10-20 stools per day was diarrhea, and anything over that was dysentery. The decision in this article was to accept as dysentery only those cases in which an etiologic agent was actually identified at O/H.

Table 3 also lists the diagnoses of malaria, based on positive antibody titers, as determined by the Center for Disease Control in Atlanta. There were four RPWs with *Plasmodium* (*P.*) *falciparum*, and eight with *P. vivax*. There were nine RPWs with mixed infection: four with *P. falciparum* + *P. vivax*, and five with three species (*P. falciparum* + *P. vivax* + *P. malariae*). Although all RPWs were given standard end-of-tour malaria prophylaxis, three had recurrences of acute attacks after discharge from the sick list; one of the attacks responded only to treatment with an experimental drug.

During captivity, 34 additional RPWs were given "quinine" for febrile episodes diagnosed by Vietnamese paramedics, without benefit of blood smears, as "malaria." These individuals were seronegative for malaria antibody titer at O/H.

Discussion

The health of the RPWs at O/H was better than had been anticipated under "worst case" planning assumptions. Nevertheless, there are a number of individuals with permanent impairments, and many others who are "at risk" because of the extended period of time during which they were subjected to protein-calorie and vitamin-malnutrition.^{5,7-9}

Eleven men reported their lowest estimated captivity weight as being 100 pounds or less. Table 3 lists the diagnoses related to malnutrition taken from the hospital records. The accuracy of these diagnoses is open to question due to several problems: first, they are based, in part, on a layman's description and memory of events which occurred several years earlier. Secondly, many of the so-called "classic" signs of vitamin deficiency are subjective (e.g., night blindness) or can have several causes (e.g., cheilosis).

Thirty-one (40 per cent) of the RPWs reported episodes of "edema" and "swelling" during captivity, including the 12 individuals with symptoms persisting up to the time of O/H (Table 1). In at least some instances, these resemble textbook descriptions of wet beriberi (pedal and ankle edema, rapidly progressing to involve upper legs and scrotum). In some cases, the Vietnamese paramedic made a diagnosis of "beriberi" and treated it with "thiamine," with

a rapid response. The one Army physician held as a POW felt, however, that these cases represented "hunger edema" due to low serum protein levels.

The signs and symptoms of clinical thiamine deficiency in the RPWs could also be attributed to traumatic peripheral nerve injury resulting from the extensive use of various restraining devices by their captors. The use of ropes, ratchet hand cuffs, leg irons, and stocks has been described in detail elsewhere.^{1-2,4,6} These often constricted an extremity, producing ischemic injury, swelling, and paresthesias, as did the forced maintenance of one position for a prolonged period. It is also possible that the swelling and paresthesias were due to an interaction between low thiamine levels and ischemic injury.

The deficiency anemias (Table 3), due to inadequate iron and/or folate, were diagnosed on the basis of serum measurements of these nutrients. The one diagnosis of "probable tropical sprue" was based upon tests indicating a malabsorption state and the appearance of the small bowel series. No small bowel biopsy was performed.

Tuberculosis was one of the three most common causes of death among RPWs in the period immediately after World War II.⁹ In contrast, there were no diagnoses of active tuberculosis among the current group of RPWs, although four men had positive PPD tests. In all of these cases, the most recent skin tests prior to captivity were negative.

There were 113 serious injuries (including fractures,

wounds, and burns, sometimes of a multiple nature). In nearly all instances, these were received in the fire fight immediately preceding capture. The majority of these injuries healed uneventfully, in spite of the minimal care received, leaving only a scar, although 11 RPWs had residual shrapnel at repatriation. One exception was open wounds of the ear. These, allegedly, were due to a captor practice of striking POWs with a cupped hand over the ear, in an attempt to break the tympanic membrane. Out of six RPWs reporting this assault, all had scarred tympanic membranes on examination.

Fourteen RPWs (18 per cent) had radiological evidence of vertebral compression fractures, an incidence level which is more characteristic of a group of aviators who had been forced to eject from jet aircraft¹ than it is of "typical" Army injuries. It may be speculated that some of these occurred during crashes of helicopters but, in point of fact, this accounts for only one case.

In spite of the fact that most of the 14 other fractures (noncompression) were secondary to gunshot wounds and the adverse conditions under which the RPWs were held, there were only two cases of osteomyelitis.

Another curious fact is the high incidence of impaired hearing, (31 RPWs or 40 per cent). Although such a finding is expected among aviators continually exposed to jet noise, why this should also be so among Army RPWs is not clear. Indeed, only 10 of the diagnoses of impaired hearing occurred among the 25 Army RPWs who were involved with helicopters or fixed wing aircraft.

Other reports have indicated that skin diseases were the greatest medical cause of time lost by soldiers fighting in Vietnam, and the POWs shared this problem. Twenty-one of the RPWs (27 per cent) had bacterial skin infections, usually boils, carbuncles, pyodermas, and cellulitis. Dermatophytoses were diagnosed in 30 of the RPWs (39 per cent); the 30 individuals represent 42 diagnoses, of which 23 were fungal foot infections.

Table 3 presents the infectious and parasitic diseases diagnosed in the RPWs. The incidence of both the helminthiases and the malarial diseases reflect the intimate sojourn of the POWs in the SVN jungles. Although the number of cases is relatively small, only among Army RPWs were any diagnoses of Schistosomiasis, Clonorchiasis, Fasciolopsiasis, and Strongyloidiasis made.

Malaria, until recently, was responsible for more deaths per year, on a world-wide basis, than any other transmissible disease. It is definitely endemic in SVN and many RPWs described what they considered to be repeated attacks of malaria, of gradually attenuating severity.

"Premature aging" is a concept frequently invoked in discussions of former POWs.⁹⁻¹¹ Evidence bearing on this matter in Vietnam RPWs is scant. Nevertheless, eight of the Army RPWs were 19 or 20 years old at time of capture, and late adolescence is still a vulnerable time to undergo the stress of malnutrition. Two RPWs presented with radiologic osteoporosis. In one RPW, this condition was restricted to the bones of the lower extremities, and was apparently due to disuse secondary to paraplegia; for the second RPW a mild osteoporosis was detected only in the lumbosacral spine and, while premature aging cannot be ruled out definitively, there was strong evidence of nutritional defi-

Table 3

Infectious and Parasitic Diseases in Army Vietnam RPWs

ICDA*	Diagnosis	Freq.	Percent
Intestinal Bacterial Infections			
003	Salmonella	2	3
004	Bacillary Dysentery	4	5
Intestinal Protozoal Infections			
40001**	E. histolytica	6	8
40002**	E. coli	12	16
40003**	E. nana	5	6
007	Giardia	2	3
40009**	Other (E. buetschilli)	1	1
40009**	Other (E. hartmani)	1	1
Malaria			
084.0	Plasmodium falciparum	13	17
084.1	Plasmodium vivax	17	22
084.2	Plasmodium malariae	5	6
Helminthiases			
120	Schistosomiasis	1	1
121.1	Clonorchiasis	1	1
121.4	Fasciolopsiasis	1	1
126	Ancylostomiasis	35	45
127.0	Ascariasis	33	43
127.1	Strongyloidiasis	3	4
127.2	Trichuriasis	24	31
127.3	Enterobiasis	7	9

* ICDA: Code numbers taken from U.S. National Center for Health International Classification of Diseases for Use in the U.S., 8th Rev.

** Code numbers created by CPMS Medical Specialties Branch

ciencies in this individual. Another interesting question is whether expected or "normal" age-related changes will occur earlier, among these POWs, than they might have otherwise. Seven RPWs (nine per cent) had degenerative changes of the spine ("spondylitis osteoarthritis") as evidenced by x-ray. There was no evidence of endocrinologic aging, such as decreased glucose tolerance.

The World War II and Korean POW post-captivity experiences were characterized by prolonged and recurrent medical and psychiatric problems.⁸⁻¹³ The precise role of the captivity experience in the genesis of those problems of earlier POWs remains controversial and undetermined. Nevertheless, increased morbidity and mortality certainly seems to exist among them.⁹ Perhaps a more favorable outcome can be expected in this new group of RPWs because of the thoroughness of the medical and dental treatment during O.H. and the planned continuation of the program of examination and treatment in future years.

References

- ¹Berg, S. W. and Richlin, M.: Injuries and illnesses of Vietnam War POWs I: Navy POWs. *Milit. Med.*, 142:515-518, 1977.
- ²Rowe, J. N.: *Five Years to Freedom*. Little, Brown, and Co., Boston, 1971.
- ³Kushner, F. H.: To live or die. *Spectrum*, 1:16-21, 1974.
- ⁴Berg, S. W.: Medical Aspects of Captivity and Repatriation. In *Family Separation and Reunion*. McCubbin, H. L., Dahl, B. B., Metres, P. J., Hunter, E. J., and Plag, J. A. (Eds.), Washington, D.C., Superintendent of Documents, 1974.
- ⁵Hill, T. M.: Dietary intake of American POWs in Vietnam. Paper presented at California Dietetic Assoc. Spring Convention, 2 May 1974.
- ⁶Rowan, S. A.: *They Wouldn't Let Us Die*. Jonathan David, Middle Village, N.Y., 1973.
- ⁷Crawford, J. N. and Reid, J. A. G.: Nutritional disease affecting Canadian troops held prisoner of war by the Japanese. *Can. J. Res.*, 25:53-85, 1947.
- ⁸Cohen, B. M. and Cooper, M. Z.: *A Follow-up Study of World War II Prisoners of War*. Veterans Admin. Med. Monograph, US Government Printing Office, Washington, D.C., 1954.
- ⁹Beebe, G. W.: Follow-up Studies of World War II and Korean War prisoners: II. Morbidity, Disability, and Maladjustments. *Am. J. Epidemiol.*, 101:400-422, 1975.
- ¹⁰Eitinger, L. and Strom, A.: *Mortality and Morbidity after Excessive Stress*. Humanities Press, New York, 1973.
- ¹¹Segal, J.: Long-term psychological and physical effects of the POW experience: A review of the literature. Technical Report 74-2, Naval Health Research Center, San Diego, 1974.
- ¹²Archibald, H. C. and Tuddenham, R. D.: Persistent stress reaction after combat. *Arch. Gen. Psychiatry*, 12:475-481, 1965.
- ¹³Nefzger, M. D.: Follow-up studies of World War II and Korean War prisoners. I. Study Plan and Mortality Findings. *Am. J. Epidemiol.*, 91:123-138, 1970.



Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 14 75-82	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) 6 Injuries and Illnesses of Vietnam War POWs. II. Army POWs.	5. TYPE OF REPORT & PERIOD COVERED 9 Final Report	
7. AUTHOR(s) 10 S. William Berg LCDR, MC, USNR Milton Richlin Ph.D.	6. PERFORMING ORG. REPORT NUMBER #	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Prisoner of War Studies Naval Health Research Center Dept. of the Navy, San Diego, CA 92152	8. CONTRACT OR GRANT NUMBER(s) 11 177C	
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research & Development Command Bethesda, Maryland 20014	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Navy N00018-75-WR-0004 Army 75-01	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Bureau of Medicine and Surgery Department of the Navy Washington, DC 20372 12 8p.	12. REPORT DATE Sept 1975	
	13. NUMBER OF PAGES	
	15. SECURITY CLASS. (of this report) unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Prisoners of War; Southeast Asia; Repatriation; Operation Homecoming; Medical Exam; Malnutrition; Infectious & Parasitic Diseases; Nerve Injury; Torture; U.S. Army.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Seventy-seven Army POWs were repatriated in Feb - March, 1973, spending up to eight years in South and North Vietnam prisons (mean duration of captivity = 49.3 months). There were 28 officers (mean age at time of capture = 27.6 years) and 49 enlisted men (mean age = 23.0 years). This report documents the injuries and illnesses diagnosed at repatriation. Although the repatriated POWs (RPWs) were generally healthier than had been expected they had an average of 14.9 diagnoses, comprising 386 separate diagnostic entities. The		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

391442

JP

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

six most prevalent diagnoses were: Helminthiases (77% of the RPWs), Avitaminoses and Other Nutritional Deficiencies (55%), Hearing Impairment (40%), Dermatophytosis (39%), Peripheral Nerve Injury (39%), and Malaria (34%). All diagnoses occurring in at least five percent of the RPWs are reported. In addition, less frequently occurring, but clinically significant diagnoses are discussed.

AF	100	101
102	103	104
105	106	107
108	109	110
111	112	113
114	115	116
117	118	119
120	121	122
123	124	125
126	127	128
129	130	131
132	133	134
135	136	137
138	139	140
141	142	143
144	145	146
147	148	149
150	151	152
153	154	155
156	157	158
159	160	161
162	163	164
165	166	167
168	169	170
171	172	173
174	175	176
177	178	179
180	181	182
183	184	185
186	187	188
189	190	191
192	193	194
195	196	197
198	199	200
201	202	203
204	205	206
207	208	209
210	211	212
213	214	215
216	217	218
219	220	221
222	223	224
225	226	227
228	229	230
231	232	233
234	235	236
237	238	239
240	241	242
243	244	245
246	247	248
249	250	251
252	253	254
255	256	257
258	259	260
261	262	263
264	265	266
267	268	269
270	271	272
273	274	275
276	277	278
279	280	281
282	283	284
285	286	287
288	289	290
291	292	293
294	295	296
297	298	299
300	301	302
303	304	305
306	307	308
309	310	311
312	313	314
315	316	317
318	319	320
321	322	323
324	325	326
327	328	329
330	331	332
333	334	335
336	337	338
339	340	341
342	343	344
345	346	347
348	349	350
351	352	353
354	355	356
357	358	359
360	361	362
363	364	365
366	367	368
369	370	371
372	373	374
375	376	377
378	379	380
381	382	383
384	385	386
387	388	389
390	391	392
393	394	395
396	397	398
399	400	401
402	403	404
405	406	407
408	409	410
411	412	413
414	415	416
417	418	419
420	421	422
423	424	425
426	427	428
429	430	431
432	433	434
435	436	437
438	439	440
441	442	443
444	445	446
447	448	449
450	451	452
453	454	455
456	457	458
459	460	461
462	463	464
465	466	467
468	469	470
471	472	473
474	475	476
477	478	479
480	481	482
483	484	485
486	487	488
489	490	491
492	493	494
495	496	497
498	499	500
501	502	503
504	505	506
507	508	509
510	511	512
513	514	515
516	517	518
519	520	521
522	523	524
525	526	527
528	529	530
531	532	533
534	535	536
537	538	539
540	541	542
543	544	545
546	547	548
549	550	551
552	553	554
555	556	557
558	559	560
561	562	563
564	565	566
567	568	569
570	571	572
573	574	575
576	577	578
579	580	581
582	583	584
585	586	587
588	589	590
591	592	593
594	595	596
597	598	599
600	601	602
603	604	605
606	607	608
609	610	611
612	613	614
615	616	617
618	619	620
621	622	623
624	625	626
627	628	629
630	631	632
633	634	635
636	637	638
639	640	641
642	643	644
645	646	647
648	649	650
651	652	653
654	655	656
657	658	659
660	661	662
663	664	665
666	667	668
669	670	671
672	673	674
675	676	677
678	679	680
681	682	683
684	685	686
687	688	689
690	691	692
693	694	695
696	697	698
699	700	701
702	703	704
705	706	707
708	709	710
711	712	713
714	715	716
717	718	719
720	721	722
723	724	725
726	727	728
729	730	731
732	733	734
735	736	737
738	739	740
741	742	743
744	745	746
747	748	749
750	751	752
753	754	755
756	757	758
759	760	761
762	763	764
765	766	767
768	769	770
771	772	773
774	775	776
777	778	779
780	781	782
783	784	785
786	787	788
789	790	791
792	793	794
795	796	797
798	799	800
801	802	803
804	805	806
807	808	809
810	811	812
813	814	815
816	817	818
819	820	821
822	823	824
825	826	827
828	829	830
831	832	833
834	835	836
837	838	839
840	841	842
843	844	845
846	847	848
849	850	851
852	853	854
855	856	857
858	859	860
861	862	863
864	865	866
867	868	869
870	871	872
873	874	875
876	877	878
879	880	881
882	883	884
885	886	887
888	889	890
891	892	893
894	895	896
897	898	899
900	901	902
903	904	905
906	907	908
909	910	911
912	913	914
915	916	917
918	919	920
921	922	923
924	925	926
927	928	929
930	931	932
933	934	935
936	937	938
939	940	941
942	943	944
945	946	947
948	949	950
951	952	953
954	955	956
957	958	959
960	961	962
963	964	965
966	967	968
969	970	971
972	973	974
975	976	977
978	979	980
981	982	983
984	985	986
987	988	989
990	991	992
993	994	995
996	997	998
999	1000	1001

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)